



ExoBacteria™ OMV Isolation Kit (for *E.coli* and other gram-negative bacteria)

Cat # EXOBAC100A-1

User Manual

Storage: Please see individual components

Version 1
3/14/2018

A limited-use label license covers this product. By use of this product, you accept the terms and conditions outlined in the License and Warranty Statement contained in this user manual.

Contents

| | |
|---|---|
| Product Description..... | 2 |
| List of Components..... | 2 |
| Storage..... | 2 |
| General Information..... | 2 |
| Protocol for ExoBacteria OMV Isolation Kit: | 3 |
| Example Data and Applications..... | 4 |
| References..... | 7 |
| Technical Support..... | 8 |
| Licensing and Warranty Statement..... | 8 |

Product Description

Bacteria-derived outer membrane vesicles (OMVs for short) have recently gained attention as a novel mechanism for transmission of molecular information within bacteria as well as bacteria to mammalian cells. Owing to its similarities to well-established extracellular vesicles (EVs), researchers have utilized OMVs to study host-bacterial interactions in pathogenesis¹, in cancer therapy², in modulating host immune response³, and engineering for use as vaccines⁴.

To assist researchers in identifying new and exciting applications for OMVs, SBI is proud to introduce the ExoBacteria™ OMV Isolation Kit – the first isolation system dedicated for OMV isolation (from *E.coli* and other Gram-negative species, e.g. *P. putida*) for use in downstream applications. Using an innovative precipitation-free, ion-exchange chromatography system (containing a proprietary capture resin + gravity column) for capture of OMVs from bacterial culture medium, the kit delivers OMVs in less than 1 hr of total time with purity and yield rivaling ultracentrifugation-based approaches.

The kits are provided with enough reagents and consumables to process up to 20 reactions*, allowing researchers maximal flexibility with experimental conditions.

***1 reaction is defined as 1 ml of resin + 30 ml of purified bacterial culture media loaded into a single gravity column**

List of Components

| Item | Volume/Qty | Storage Temperature |
|----------------------|------------|---------------------|
| OMV Binding Resin | 20 mL | 4°C |
| Binding Buffer (20X) | 60 mL | 4°C |
| Elution Buffer (1X) | 30 mL | RT |
| Gravity Flow Columns | 20 | RT |
| Column Stoppers | 20 | RT |
| Column Caps | 20 | RT |

Storage

The Kit is shipped on Blue Ice and the components should be stored at recommended temperatures as stated above. Properly stored kits are stable for 12 months from the date received.

General Information

Note: The Binding Buffer is provided in a 20X concentration, please dilute to a 1X working concentration in DI water prior to starting the procedure.

Materials required but not provided: Filtering units with 0.45 µm and 0.22 µm filters, column rack, rotating rack, pipettes and pipette tips

Protocol for ExoBacteria OMV Isolation Kit:

1. Prepare clarified supernatant from bacterial culture

- a. Culture the bacteria in its appropriate condition until the desired OD is reached.
- b. Spin down bacteria at 5000 x g for 20 mins at 4°C.
- c. Transfer the supernatant to a fresh tube and spin again at 5000 x g for 20 mins at 4°C to completely remove any cellular debris.
- d. Collect the culture supernatant and filter through a 0.45 µm vacuum filter.
- e. Repeat the filtration step with a 0.22 µm vacuum filter and collect the material.

Filtered culture supernatant is now ready for OMV isolation.

2. Column preparation and OMV capture

- a. Mix the OMV Binding Resin by shaking the bottle thoroughly as contents may have settled at the bottom
- b. Pipette 1 mL of the OMV Binding Resin onto the column.
- c. Equilibrate by adding 10 mL of the 1X Binding Buffer to column, snap off the existing column tip cap, and allow solution to completely flow through. Discard the flow through.
- d. Place a yellow column stopper onto the bottom of the column.
- e. Add 30 mL of the clarified bacterial culture supernatant (prepared in step 1) to the resin and place column cap on top of the column.
- f. Incubate on a rotating rack at 4°C for 30 minutes to allow for OMV binding

3. OMV Elution

- a. Place the column onto a rack and uncap the top and bottom of the column
- b. Allow the resin/supernatant mixture to flow through
- c. Wash the resin with 15 mL Binding Buffer. Discard the flow through. Repeat the wash step 2x.
- d. Cap the column at the bottom with the stopper and add 1.5 mL OMV Elution Buffer. Incubate for 2 mins at room temperature, gently shaking the column every 30 secs.
- e. Uncap the column at the bottom and collect the eluate containing the OMVs in a fresh microcentrifuge tube.

Example Data and Applications

A. ExoBacteria OMV Isolation Kit

B. Ultracentrifugation





Figure 1. OMVs isolated using the ExoBacteria OMV Isolation Kit are similar in appearance, but with tighter distribution than OMVs isolated via ultracentrifugation. Comparison of transmission electron micrographs (TEM) of *E. coli*-derived OMVs isolated using (A) the ExoBacteria OMV Isolation Kit or (B) ultracentrifugation. OMVs (light blue arrows) are indicated in both samples, but note the unwanted protein aggregates (red arrow) that appear in the ultracentrifugation sample.

A. Particle size distribution



B. Yield

| | OMV Yield (particles/ μg protein) |
|-------------------------------|---|
| ExoBacteria OMV Isolation Kit | 1.38×10^8 |
| Ultracentrifugation | 0.0657×10^8 |

Figure 2. The ExoBacteria OMV Isolation Kit delivers a narrower size distribution of OMVs and higher yields than ultracentrifugation. (A) Comparison of fluorescent NTA analysis of *E. coli*-derived bacterial OMVs isolated using either ExoBacteria OMV Isolation Kit (blue line) or ultracentrifugation (gray line) shows how the ExoBacteria OMV Isolation Kit delivers a much narrower size distribution of isolated OMVs than conventional ultracentrifugation approach. (B) In addition, total yield is 20-fold higher with the ExoBacteria OMV Isolation Kit (1.38×10^8 particles/ μg of protein) compared to ultracentrifugation (0.0657×10^8 particles/ μg of protein).

A. Detection of OMV marker



B. Detection of protein carry-over





Figure 3. Get higher OMV yield and less carry-over protein with the ExoBacteria OMV Isolation Kit. Western blot analyses of *E. coli*-derived OMVs isolated using either the ExoBacteria OMV Isolation Kit or ultracentrifugation show that **(A)** the ExoBacteria OMV Isolation Kit delivers higher yield of OMVs than ultracentrifugation—compare the amounts of GroEL in each lane—and **(B)** the ExoBacteria OMV Isolation Kit preps have less carry-over protein than the ultracentrifugation preps—note the lower amount of flagellin. Each lane is loaded with 5 μ g of total protein.

A. Distinct proteins identified by mass spectrometry



B. Number of OMV proteins found in EV-Pedia




Figure 4. Identify more OMV proteins with OMVs isolated using the ExoBacteria OMV Isolation Kit. (A) Comparison of the total number of proteins identified via mass spectrometry from OMV samples isolated using either the ExoBacteria OMV Isolation Kit or ultracentrifugation shows a wider range of proteins identified from OMVs isolated with the ExoBacteria OMV Isolation Kit. All but 7 of the proteins identified in the OMVs isolated using ultracentrifugation were also found in the OMVs isolated using the ExoBacteria kit, suggesting that the higher yield and higher purity delivered by the ExoBacteria OMV Isolation Kit expand your ability to derive insights from OMVs. **(B)** More of the proteins from the ExoBacteria OMV kit-isolated OMVs were found in the EV-Pedia database than from OMVs isolated using ultracentrifugation, demonstrating the increased specificity of the ExoBacteria OMV Isolation Kit.

References

1. Vanaja SK *et al.* Bacterial Outer Membrane Vesicles Mediate Cytosolic Localization of LPS and Caspase-11 Activation. *Cell* 2016 May 19; 165(5): 1106-1119.
2. Kim OY *et al.* Bacterial outer membrane vesicles suppress tumor by interferon- γ -mediated antitumor response. *Nat Commun.* 2017 Sep 20;8(1):626. doi: 10.1038/s41467-017-00729-8

3. Vidakovics ML *et al.* B cell activation by outer membrane vesicles--a novel virulence mechanism. PLoS Pathog. 2010 Jan 15;6(1):e1000724. doi: 10.1371/journal.ppat.1000724.

4. Chen DJ *et al.* Delivery of foreign antigens by engineered outer membrane vesicle vaccines. Proc Natl Acad Sci U S A. 2010 Feb 16;107(7):3099-104. doi: 10.1073/pnas.0805532107. Epub 2010 Jan 27.

Technical Support

For more information about SBI products and to download manuals in PDF format, please visit our web site: <http://www.systembio.com>

For additional information or technical assistance, please call or email us at:

System Biosciences (SBI)
2438 Embarcadero Way
Palo Alto, CA 94303

Phone: (650) 968-2200
Toll-Free: (888) 266-5066
Fax (650) 968-2277

E-mail:

General Information: info@systembio.com
Technical Support: tech@systembio.com
Ordering Information: orders@systembio.com

Licensing and Warranty Statement

Limited Use License

Use of the ExoBacteria OMV Isolation Kit (*i.e.*, the "Product") is subject to the following terms and conditions. If the terms and conditions are not acceptable, return all components of the Product to System Biosciences (SBI) within 7 calendar days. Purchase and use of any part of the Product constitutes acceptance of the above terms.

The purchaser of the Product is granted a limited license to use the Product under the following terms and conditions:

- The Product shall be used by the purchaser for internal research purposes only. The Product is expressly not designed, intended, or warranted for use in humans or for therapeutic or diagnostic use.
- The Product may not be resold, modified for resale, or used to manufacture commercial products without prior written consent of SBI.
- This Product should be used in accordance with the NIH guidelines developed for recombinant DNA and genetic research.

SBI has pending patent applications related to the Product. For information concerning licenses for commercial use, contact SBI.

Purchase of the product does not grant any rights or license for use other than those explicitly listed in this Licensing and Warranty Statement. Use of the Product for any use other than described expressly herein may be covered by patents or subject to rights other than those mentioned. SBI disclaims any and all responsibility for injury or damage which may be

caused by the failure of the buyer or any other person to use the Product in accordance with the terms and conditions outlined herein.

Limited Warranty

SBI warrants that the Product meets the specifications described in this manual. If it is proven to the satisfaction of SBI that the Product fails to meet these specifications, SBI will replace the Product or provide the purchaser with a refund. This limited warranty shall not extend to anyone other than the original purchaser of the Product. Notice of nonconforming products must be made to SBI within 30 days of receipt of the Product.

SBI's liability is expressly limited to replacement of Product or a refund limited to the actual purchase price. SBI's liability does not extend to any damages arising from use or improper use of the Product, or losses associated with the use of additional materials or reagents. This limited warranty is the sole and exclusive warranty. SBI does not provide any other warranties of any kind, expressed or implied, including the merchantability or fitness of the Product for a particular purpose.

SBI is committed to providing our customers with high-quality products. If you should have any questions or concerns about any SBI products, please contact us at (888) 266-5066.

© 2018 System Biosciences (SBI), All Rights Reserved

This page intentionally left blank.

This page intentionally left blank.

This page intentionally left blank.



System Biosciences (SBI)
2438 Embarcadero Way
Palo Alto, CA 94303

Phone: (650) 968-2200
Toll-Free: (888) 266-5066
Fax: (650) 968-2277

E-mail:
General Information: info@systembio.com
Technical Support: tech@systembio.com
Ordering Information: orders@systembio.com