

PRODUCT	Purified Human and Mouse Exosomes
CATALOG #s	EXOP-100A-1, EXOP-105A-1, EXOP-110A-1, EXOP-115A-1, EXOP-120A-1, EXOP-125A-1, EXOP-130A-1, EXOP-135A-1, EXOP-140A-1, EXOP-145A-1, EXOP-200A-1, EXOP-300A-1
SIZE	50 ug (>1x10⁶ frozen exosomes)
LOT #	All
STORAGE	-20°C to -80°C
SHELF LIFE	12 months from date of receipt with proper storage
SHIPPING	Dry Ice

PACKAGE CONTENTS

Each purified exosome kit comes in 50 ul sterile 1x PBS with 50 ug exosomes (>1x10⁶ frozen exosomes).

Catalog#	Description	Size
EXOP-100A-1	MCF-7 Human breast cancer, noninvasive cell line: >1x10 ⁶ frozen exosomes	50 ug
EXOP-105A-1	MDA-MB-231 Human breast cancer, aggressive/invasive/metastatic cell line: >1x10 ⁶ frozen	50 ug
EXOP-110A-1	HEK293 Human embryonic kidney cell line: >1x10 ⁶ frozen exosomes	50 ug
EXOP-115A-1	PC-3 Human prostate cancer cells derived from metastatic cancer cell line: >1x10 ⁶ frozen exosomes	50 ug
EXOP-120A-1	A549 Human non-small cell lung cancer cell line: >1x10 ⁶ frozen exosomes	50 ug
EXOP-125A-1	H841 Human small cell lung cancer cell line: >1x10 ⁶ frozen exosomes	50 ug
EXOP-130A-1	H196 Human small cell lung cancer cell line: >1x10 ⁶ frozen exosomes	50 ug
EXOP-135A-1	DMS114 Human small cell lung cancer cell line: >1x10 ⁶ frozen exosomes	50 ug
EXOP-140A-1	PCS-500-011 Human pre-adipose derived mesenchymal stem cell: >1x10 ⁶ frozen exosomes	50 ug
EXOP-145A-1	PCS-500-012 Human bone marrow-derived mesenchymal stem cell line: >1x10 ⁶ frozen exosomes	50 ug
EXOP-200A-1	JAWSII Mouse bone marrow immature dendritic cell line: >1x10 ⁶ frozen exosomes	50 ug
EXOP-300A-1	Human pooled serum: >1x10 ⁶ frozen exosomes	50 ug


DESCRIPTION

Exosomes are 60 - 180 nm membrane vesicles secreted by most cell types in vivo and in vitro and contain distinct subsets of RNAs and proteins depending upon the cell type from which they are secreted, making them useful for biomarker discovery and functional characterization. Exosomes are nano-sized shuttles that transport signaling RNAs, lipids and proteins to other cells. Studying exosome contents are thus a "liquid biopsy" for biomarkers to gain insights into their roles in disease initiation and progression.

- **Cell lines grown in exosome-depleted FBS (Exo-FBS)**
- **Exosomes purified using ExoQuick-TC**
- **Characterized by NanoSight for size and intactness**
- **CD63 Western blot analysis validated**
- **Exosomes at >1x10⁶ exosomes (50 ug protein)**

BIOMARKER VALIDATION

Western blot on purified exosomes




#	Line	Type
1	MCF-7	Human breast cancer, noninvasive cell line exosomes
2	MDA-MB-231	Human breast cancer, aggressive/invasive/metastatic cell line exosomes
3	PC-3	Human prostate cancer cells derived from metastatic cancer cell line exosomes
4	A549	Human non-small cell lung cancer cell line exosomes
5	H196	Human small cell lung cancer cell line exosomes
6	H841	Human small cell lung cancer cell line exosomes
7	HEK293	Human embryonic kidney cell line exosomes
8	DMS114	Human small cell lung cancer cell line exosomes
9	PCS-500-011	Human pre-adipose derived mesenchymal stem cell exosomes
10	PCS-500-012	Human bone marrow-derived mesenchymal stem cell line exosomes
11	JAWSII	Mouse bone marrow immature dendritic cell line exosomes
12	Human Serum	Human pooled serum healthy exosomes

An aliquot of the purified exosomes from the cell lines and from human serum were lysed with RIPA buffer to make exosomes protein lysates. Approximately 20 ug of protein for each sample was separated on a gradient SDS-PAGE and then transferred to nitrocellulose membranes. The membranes were probed for CD63 profiles using SBI's anti-CD63 antibody (cat# EXOAB-CD63A-1) at a 1:1,000 dilution. Bands were detected using the secondary HRP-conjugated antibody at 1:10,000 and blots imaged. All purified exosomes preparations are positive, immunoreactive for CD63 and produces variable banding patterns common to published exosome CD63 profiles.

NANOSIGHT DATA

Approximately 5 ul of the purified exosomes sample was added to 995 ul of 0.2 um filtered 1X PBS (1:200 dilution). The diluted samples were incubated in a VWR 500 model ultrasonicator water bath set at 33°C for 10 minutes to ensure adequate exosome particle dispersion. The samples were diluted 1:10 then vortexed at 2.5k for 10 seconds. This eventual 1:2,000 dilution was used to gather between 1,000 to 3,000 particle tracks per sample analysis. The samples were then loaded into a NanoSight LM10HSB with a syringe pump and the sensitivity of the camera is set to auto 16 (the most sensitive auto-setting). All data were collected in triplicate. The purified exosomes displayed the expected size distribution profiles, with peak diameters between 90-110 nm and concentrations in the range expected for media exosomes at about 1x10¹⁰ exosomes/ml.



The Product shall be used by the purchaser for internal research purposes only. The Product is expressly not designed, intended, or warranted for use in humans or for therapeutic or diagnostic use. This Product is covered by a limited-use label license. Purchase and use of any part of the Product constitutes acceptance of the terms outlined in the Licensing and Warranty Statement contained in the User Manual.