



EVeryRNA™ EV RNA Purification System with ExoQuick-TC EV Isolation

Cat # EVery106TC-1
User Manual

Storage:

DNase I and Glycogen at -20°C
All other components at room temperature

Version 2
3/17/2022

A limited-use label license covers this product. By use of this product, you accept the terms and conditions outlined in the License and Warranty Statement contained in this user manual.

Contents

Product Description	2
List of Components	4
Additional Required and Optional Equipment Not Included in Kit	4
Protocol	5
Example Data and Applications	7
Technical Support	11
Licensing and Warranty Statement	11

Product Description

Get easy EV isolation with ExoQuick-TC and total EV RNA purification with EVeryRNA in one convenient kit—high yields and high-quality drive reliable insight.

Discover more when you capture total EV RNA. Combine the time- and labor-savings of ExoQuick-TC EV Isolation technology with the total RNA isolation capabilities of EVeryRNA EV RNA Purification technology for a complete solution for your EV RNA biomarker workflows.

The EVeryRNA EV RNA Purification System overcomes many of the challenges faced when isolating RNA from extracellular vesicles (EVs), most notable is the ability to capture total EV RNA, including small RNAs. EVeryRNA is effective even with low amounts of input RNA and is capable of delivering high yields of highly pure RNA. Because the RNA elutes in a small sample volume, thus generating a highly concentrated prep, you can increase the amount of RNA used in a single downstream reaction for better data coverage quality.

- Move quickly and confidently with exoRNA isolation that's high-yield and complete in <30 minutes
- Find what others miss when you capture every RNA with EVeryRNA
- Achieve phenol-level yields with a safer column-based method
- Get more RNA for each downstream reaction with EVeryRNA's small-volume elutions
- Ensure delivery of highly pure RNA by using the included DNaseI
- Isolate RNA from EVs for a full range of downstream applications, such as RNA-seq and miRNA profiling
- Maximize productivity with ExoQuick-TC EV Isolation bundled with EVeryRNA

The EVeryRNA EV RNA Purification System with ExoQuick-TC EV Isolation comes with sufficient reagents to perform 10 EV isolation reactions from tissue culture medium or other biofluids and then EV RNA purification. EVeryRNA technology is available bundled with several of SBI's powerful EV isolation technologies as well as a cDNA Synthesis and Pre-amplification Kit (Table 1).

Table 1. EVeryRNA EV RNA Purification Products

Catalog number	Description
EVery100A-1	EVERYRNA™ EV RNA Purification System
EVery106EQ-1	EVERYRNA™ EV RNA Purification System with ExoQuick EV Isolation
EVery106TC-1	EVERYRNA™ EV RNA Purification System with ExoQuick-TC EV Isolation
EVery106SS-1	EVERYRNA™ EV RNA Purification System with SmartSEC Single
EVery200A-1	EVERYRNA™ cDNA Synthesis & Pre-Amplification Kit
EVery300A-1	EVERYRNA™ EV RNA Purification System & cDNA Synthesis Kit (includes EVery100A-1 and EVery200A-1)

Workflow: Maximize productivity with ExoQuick-TC and EVeryRNA

Step 1: Isolate EVs with ExoQuick-TC

Isolating EVs from tissue culture media, saliva, urine, follicular fluid, breast milk, and other biofluids with ExoQuick-TC is quick and easy, with a simple workflow that requires only 10-minutes of hands-on time (Figure 1).

To isolate exosomes from tissue culture media, simply:

- Add an appropriate volume of ExoQuick-TC
- Incubate overnight at 4°C
- Isolate exosomes with a 30-minute low-speed spin (1500g).

Isolated exosomes can be found in the pellet and resuspended in an appropriate solution.




Figure 1. The ExoQuick-TC workflow.

Step 2: Purify EV RNA with EVeryRNA

The EVeryRNA EV RNA Purification System delivers high yields of highly concentrated RNA from already isolated EVs. The column-based workflow is easy to implement and can be completed in less than 30 minutes (Figure 1).




Figure 2. The quick and easy EVeryRNA EV RNA purification workflow.

List of Components

Table 2. Components of EVery106TC-1, EVeryRNA™ EV RNA Purification System

Components	Qty/Volume	Storage Temperature
ExoQuick-TC	10 mL	RT
Lysis buffer	12.5 mL	RT
Lysis additive	1.5 mL	RT
Glycogen	50 µL	-20°C
Wash solution	18 mL	RT
Elution solution	0.5 mL	RT
Micro spin column	10	RT
Collection tubes	10	RT
Elution Tubes	10	RT
RNase free DNase I	100 µL	-20°C
Enzyme buffer	1 mL	RT

Note: The table above is for 10 reaction kit.

Additional Required and Optional Equipment Not Included in Kit

1. 96-100% Ethanol
2. β-mercaptoethanol (cat# M3148-25ML, Sigma)- optional, but highly recommended


Protocol

Step 1: Isolate EVs with ExoQuick-TC

Note: This protocol is a starting point for most cells in culture. Different cell types secrete different amount of exosomes. You may need to scale this protocol accordingly.

1. Collect biofluid and centrifuge at $3000 \times g$ for 15 minutes to remove cells and cell debris.
2. Transfer supernatant to a sterile vessel and add the appropriate volume of ExoQuick-TC to the bio-fluid. Mix well by inverting or flicking the tube.
3. Refrigerate overnight (at least 12 hours) at $+4^{\circ}\text{C}$. The tubes should not be rotated or mixed during the incubation period and should remain upright.
4. Centrifuge ExoQuick-TC/biofluid mixture at $1500 \times g$ for 30 minutes. Centrifugation may be performed at either room temperature or $+4^{\circ}\text{C}$ with similar results. After centrifugation, the exosomes may appear as a beige or white pellet at the bottom of the vessel.

Exosome pellets obtained from 10 ml of cerebral spinal fluid using ExoQuick.



5. Aspirate supernatant. Spin down residual ExoQuick-TC solution by centrifugation at $1500 \times g$ for 5 minutes. Remove all traces of fluid by aspiration, taking great care not to disturb the precipitated exosomes in pellet.
6. Resuspend exosome pellet in 100-500 μL using sterile 1X PBS, or **specific buffer according to your downstream application**. We recommend using the precipitated exosomes immediately rather than freezing them for future use.

Step 2: Purify EV RNA with EVeryRNA

Before you start the protocol for exosomal RNA isolation:

1. The protocol is outlined for 500 μL of sample input. The pellet from **ExoQuick TC** should be resuspended in 1xPBS to 500 μL to prevent column clogging during isolation.
2. All steps should be performed at room temperature and all centrifugation steps performed at $\geq 8000 \times g$ ($\geq 10,000$ rpm for table top microcentrifuge).

3. It is highly recommended to warm up **Lysis Buffer** at 60°C for 5-10 minutes and mix well until the solution becomes clear again if precipitates are present.

! OPTIONAL (highly recommended)

The use of β -mercaptoethanol in the Lysis Buffer is highly recommended. Add 10 μ L of β -mercaptoethanol to each 1 mL of **Lysis buffer**.

! OPTIONAL (highly recommended)

Add 5 μ L of Glycogen to the Lysis Mix if you are expecting a low RNA yield.

4. Prepare a working solution of the **Wash solution** by adding 42 mL of 96-100% ethanol to the supplied bottle containing the concentrated Wash Solution.
5. The RNA yield may be increased by using **Elution Buffer** warmed to 60°C.

RNA isolation steps:

1. Add 700 μ L of **Lysis Buffer**, 110 μ L of **Lysis Additive** and 7 μ L of β -mercaptoethanol (optional) to the 500 μ L PBSx1 Buffer containing the purified exosomes.
2. Mix well by vortexing for 10 sec. then incubate at RT for 10 min.

! OPTIONAL (highly recommended)

Add 5 μ L of Glycogen to the lysis mix if you know that RNA yield is low.

3. After incubation add 1.35 mL of 96%-100% EtOH to the mix from Step3 and mix well by vortexing for 10 seconds.
4. Transfer 750 μ L of the mixture from Step 4 into a Micro Spin column. Centrifuge for 1 minute. Discard the flowthrough and reassemble the spin column in its collection tube.
5. Repeat Step 5 four times to transfer the remaining mixture from Step 4 into the Micro Spin column.
6. Apply 400 μ L of **Wash Solution** on the column and centrifuge for 2 minutes. Discard the flowthrough and reassemble the spin column in its collection tube.

! OPTIONAL (highly recommended)

On-column DNA removal:

- a. For every on-column reaction prepare a mix of 7.5 μ L of **RNase-free DNase I** and 50 μ L of **Enzyme buffer**. Mix gently by inverting the tube a few times or flicking the tube with your fingers to mix.
- ! DO NOT VORTEX**
- b. Apply 57 μ L of DNase I mix from step a. to the column and incubate at 25°C-30°C for 15 minutes.
7. Apply 600 μ L of **Wash solution** to the column and centrifuge for 30 seconds. Discard the flowthrough and reassemble the spin column in its collection tube.
 8. Repeat step 9 one more time, for a total of two washes.

9. Centrifuge the empty column for 1 minute to completely remove any residual solution. Discard the collection tube.
10. Transfer the spin column to a fresh Eppendorf tube. Apply 10 μL to 25 μL of **Elution solution** to the column and let it stand for 2 minutes. Centrifuge for 1 minute to elute.
11. To maximize the recovery of the RNA add the eluate collected from step 12 back on the column, let it stand for 2 minutes. Centrifuge for 1 minute to elute.
12. Exosomal RNA is now ready for downstream applications.

Example Data and Applications

Characterizing ExoQuick-TC exosomes with NanoSight

Exosomes purified with ExoQuick-TC from tissue culture media show the expected particle size distribution and high concentration yields when analyzed using NanoSight's Nanoparticle Tracking Analysis (NTA, Figure 3).




Figure 3. Exosome size distribution and yields from tissue culture media. Cells from a human HT1080 lung sarcoma cell line were cultured in conditioned media (serum-free) for 72 hours. 10 mL of media was combined with 2 mL ExoQuick-TC, incubated overnight at 4°C, and centrifuged at 1500g for 30 minutes to isolate exosomes. The exosome pellet was resuspended in 1 mL PBS, diluted 1:40, and visualized on the NanoSight LM10 instrument. The analysis shows that ExoQuick-TC recovered 133 nm exosomes at a concentration of 1.74 x 10⁹ particles/mL.

EVERyRNA captures Everything

To demonstrate the ability of the EVERyRNA EV RNA Purification System to capture the full range of RNAs, we used the kit to isolate RNA from 10,000 cells (Figure 4, lane 1), from EVs that were isolated from 250 μL of serum

using **SmartSEC Single** (Figure 4, lane 2), and from buffer spiked with 0.1 pmol of Cel-miR-39 (Figure 2, lane 3). The high quality of the isolated RNA can be seen in lane 1, where the RNA integrity number (RIN) is 9.9 and the 28S/18S RNA ratio is 1.5. The multiple bands in lane 2 demonstrate that EVeryRNA captures RNAs of different lengths—EVERYthing—from EVs with no apparent bias or size preference. The strong signal from the spiked-in miRNA in lane 3 demonstrates the good recovery of even small RNAs.



Figure 4. EVeryRNA captures EVERYthing.

EVERYRNA works well with ExoQuick, yielding broad and unbiased size distribution of RNAs (Figure 5).




Figure 5. EVeryRNA is compatible with EVs isolated using ExoQuick.

EVERYRNA delivers similar amounts of RNA as phenol-based methods

To demonstrate the excellent RNA yields and robust cDNA synthesis obtained with EVeryRNA, we isolated EVs from 250 μ L of serum using **SmartSEC Single**, spiked in 0.1 pmol of Cel-miR-39, and used both the **EVERYRNA EV Purification System** and a phenol-based kit to isolate RNA. The isolated RNA was reverse transcribed using the EVeryRNA cDNA Synthesis & Pre-amplification Kit and the copy number of Cel-miR-39 measured (Figure 6). The EVeryRNA EV Purification System delivered similar levels of Cel-miR-39 as the phenol-based method.




Figure 6. EVeryRNA is EVery bit as good as phenol.

EVERYRNA efficiently isolates mRNA

We used the EVeryRNA EV RNA Purification System and EVeryRNA cDNA Synthesis & Pre-amplification Kit to isolate mRNA and synthesize cDNA from cells overexpressing eGFP (Figure 7). Robust levels of eGFP mRNA are recovered and converted to cDNA when cells are overexpressing eGFP.




Figure 7. EVeryRNA efficiently isolates longer RNAs like mRNA

miRNA isolated and converted to cDNA using EVeryRNA can be used for miRNA profiling

We isolated EVs from 250 µL of serum using SmartSEC Single and used the EVeryRNA EV RNA Purification System and EVeryRNA cDNA Synthesis & Pre-amplification System to isolate and reverse transcribe EV RNAs for miRNA profiling using the [SeraMir Human Exosome RNA Profiling Plate](#). We were able to detect a number of miRNAs both with amplification (96 miRNAs) and without amplification (16 miRNAs, with 14 overlapping with the miRNAs detected with amplification, Figure 8).




Figure 8. EVeryRNA generates high-quality RNA suitable for miRNA profiling both with and without amplification.

We were also able to show robust, successful RNA-seq runs using RNA isolated from EVs with EVeryRNA (Table 3). All three EV isolation methods tested generated high-quality RNA-seq data.

Table 3. Successful RNA-seq with EVeryRNA-isolated EV RNA

EV Isolation Method	Amount of RNA isolated (ng)	Number of reads	FAST-QC
ExoQuick	4.3	53,349,528	Passed
ExoQuick Ultra	5.4	107,154,128	Passed
SmartSEC Single	5.2	98,886,924	Passed

Technical Support

For more information about SBI products and to download manuals in PDF format, please visit our web site: <http://www.systembio.com>

For additional information or technical assistance, please call or email us at:

System Biosciences (SBI)
2438 Embarcadero Way
Palo Alto, CA 94303

Phone: (650) 968-2200
Toll-Free: (888) 266-5066
Fax (650) 968-2277

E-mail:
General Information: info@systembio.com
Technical Support: tech@systembio.com
Ordering Information: orders@systembio.com

Licensing and Warranty Statement

Limited Use License

Use of the EVeryRNA EV RNA Purification Products (*i.e.*, the “Product”) is subject to the following terms and conditions. If the terms and conditions are not acceptable, return all components of the Product to System Biosciences (SBI) within 7 calendar days. Purchase and use of any part of the Product constitutes acceptance of the above terms.

The purchaser of the Product is granted a limited license to use the Product under the following terms and conditions:

- The Product shall be used by the purchaser for internal research purposes only. The Product is expressly not designed, intended, or warranted for use in humans or for therapeutic or diagnostic use.
- The Product may not be resold, modified for resale, or used to manufacture commercial products without prior written consent of SBI.
- This Product should be used in accordance with the NIH guidelines developed for recombinant DNA and genetic research.

This Product is covered by licensed Patents. For information concerning licenses for commercial use, contact SBI.

Purchase of the product does not grant any rights or license for use other than those explicitly listed in this Licensing and Warranty Statement. Use of the Product for any use other than described expressly herein may be covered by patents or subject to rights other than those mentioned. SBI disclaims any and all responsibility for injury or damage which may be caused by the failure of the buyer or any other person to use the Product in accordance with the terms and conditions outlined herein.

Limited Warranty

SBI warrants that the Product meets the specifications described in this manual. If it is proven to the satisfaction of SBI that the Product fails to meet these specifications, SBI will replace the Product or provide the purchaser with a refund. This limited warranty shall not extend to anyone other than the original purchaser of the Product. Notice of nonconforming products must be made to SBI within 30 days of receipt of the Product.

SBI's liability is expressly limited to replacement of Product or a refund limited to the actual purchase price. SBI's liability does not extend to any damages arising from use or improper use of the Product, or losses associated with the use of additional materials or reagents. This limited warranty is the sole and exclusive warranty. SBI does not provide any other warranties of any kind, expressed or implied, including the merchantability or fitness of the Product for a particular purpose.

SBI is committed to providing our customers with high-quality products. If you should have any questions or concerns about any SBI products, please contact us at (888) 266-5066.

© 2020 System Biosciences (SBI), All Rights Reserved

This page intentionally left blank.

This page intentionally left blank.

This page intentionally left blank.



System Biosciences (SBI)
2438 Embarcadero Way
Palo Alto, CA 94303

Phone: (650) 968-2200
Toll-Free: (888) 266-5066
Fax: (650) 968-2277

E-mail:

General Information: info@systembio.com
Technical Support: tech@systembio.com
Ordering Information: orders@systembio.com